
**WORKSHOP SUMMARY** 

#### APPENDIX E (continued)



#### **Powerpoint Presentations**




### Ecological impacts of Climate Change in Southern California

Elsa Cleland
Division of Biological Sciences
University of California, San Diego



### Global environmental changes are altering climate, biogeochemical cycles, and species composition






-Rising atmospheric CO<sub>2</sub>

-Warming

-Shifting precipitation



-Nitrogen deposition



-Invasive species, altered fire regimes


#### California: a biodiversity hotspot

- California floristic province is one of the 10 most diverse areas on earth
- 2000+ endemic plant species
- San Diego County + 1500 total plant species





#### <u>Outline</u>



- Species responses to climate change (observed/predicted)
  - Phenology
  - Range shifts (caveats)
- Coastal sage scrub
  - Case studies: checkerspot butterfly, CA gnatcatcher
  - Response of CSS vs invasive plants to altered rainfall
  - Additional factors: Fire, N deposition
  - Restoration in the face of invasion and environmental change

## Phenology (timing) is a key component of the ecology of a species


- Phenology: study of developmental events in the life of an organism
- Species are often active at different times during the growing season
- Complementary phenology enables coexistence in diverse communities



### Shifting phenology provides evidence that global change is already impacting species and ecosystems







http://modis.gsfc.nasa.gov/

-Are these changes in phenology predictive of species at risk of declining with future climate change?

Cleland et al. Trends in Ecology and Evolution 2007

## Could phenological sensitivity predict species performance responses to warming?

- Meta-analysis of warming experiments
- Paired
   observations of
   phenology and
   performance
   (biomass, seed
   production, etc.)
- Within and across sites
  - Is local monitoring worth it?



Harvard Forest Soil Warming




Jasper Ridge Global Change Experiment




Open-top chamber, White Mountains CA

### Hypothesis: Species which phenologically "track" climate should maintain performance with warming

- Most phenological sensitivities are negative (earlier flowering per ° warming)
- We expected a negative relationship between phenological sensitivity and performance (i.e. "tracking" the thermal niche is adaptive)




### Across all sites: Species that "track" climate have higher performance



## Within sites phenological sensitivity is a better predictor of performance

- Jasper Ridge Global Change Experiment (9 species)
  - Annual grassland, in the interior foothills of coast range central California
- Rocky Mountain Biological Lab Warming Meadow (7 species)
  - Alpine meadow in Colorado
- Land managers can use phenological monitoring to identify species at risk of declining with climate change




### Herbarium specimens show earlier flowering in San Diego County – signal of warming?






www.calphotos.berkeley.edu



Data source: SDMNH Herbarium specimens, San Diego Plant Atlas, Mary Ann Hawke & Layla Aerne


#### Outline



- Species responses to climate change (observed/predicted)
  - Phenology
  - Range shifts (caveats)
- Coastal sage scrub
  - Case studies: checkerspot butterfly, CA gnatcatcher
  - Response of CSS vs invasive plants to altered rainfall
  - Additional factors: Fire, N deposition
  - Restoration in the face of invasion and environmental change

Basic assumption: With increasing temperatures, we expect species to move poleward and up in elevation, also constrained by precipitation & seasonality

In recent decades, have we seen this pattern?







Fig. 3. Changing vegetation distribution from 1977 to 2006–2007. Elevation distribution of the mean normalized vegetation coverage of the ten most widely distributed species in 1977 and 2006–2007.



- -Ex. Small mammal communities in Yosemite National Park
- -2003-2006 Resampled areas originally monitored by Grinnell & colleagues 1914-1920
- -3ºC increase in temperatures over this time






Fig. 1. Map of surveyed sites in Grinnell (Historic) and Current surveys relative to the Yosemite National Park boundary and life zones (upper panel), and to an averaged elevational profile (lower panel).

Moritz et al. 2008 Science

- -Average increase in elevational ranges of small mammals
- -Species responses were mixed, some moved down
- -Much of this average shift was due to contractions at the lower edge of the elevational range









Shifts in plant species elevational range limits and abundances observed over nearly five decades in a western North America mountain range

Christopher W. Kopp & Elsa E. Cleland




- -2010 resurvey of elevational transects in the White Mountains originally collected in 1961
- -Individualistic species responses in abundance and elevational range (no average shift upwards)
- -Sagebrush moved rapidly upward and became much more abundant
- -Sagebrush encroachment potentially contributed to large declines in cushion plants
- -Conclusion: altered species interactions with climate change

#### How to predict where species will move?



#### Projected range shifts for rare species






Shaw et al 2009, California Climate Change Center Draft Report "The Impact of Climate Change on California's Ecosystem Services"

### Caveats: Species Distribution Models (SDMs)

- -Assume a set dispersal distance
- -Lots of barriers to dispersal
- -Other habitat requirements (e.g. soils, biotic interactions)
- -Velocity of climate change is faster for flat areas than steep elevation gradients, species need to disperse further faster in areas without elevational gradients



**Figure 1** | **Changing temperature in California. a**, Current (1950–2000) mean annual temperature at 800 m resolution. The black rectangle indicates the Central California inset in **b. c**, The spatial gradient of temperature change using a 9 pixel kernel. **d**, The temporal gradient of climate change from 2000–2099 from 0.5 °C 16 general circulation model (GCM) ensemble projection with A1B emissions. **e**, The velocity of climate change determined from the quotient of **d** and **c**.

-Species
Distribution Models
(SDMs) have
become more
advanced

-Ex. Network chain analysis recognizes that species will need to move across landscapes over time

-Identifies critical areas of connectivity

-Outputs of SDM predictions available





Figure 2.9: Required Chains of Salvia Leucophylla (Blue) Overlaid on All Connectivity Chains Formed Under GFDL Climate Projections Through 2080. Model outputs shown assume 1.5 pixel dispersal distance and 1,000km<sup>2</sup> suitable habitat (57 chains) requirement.

#### Outline




- Species responses to climate change (observed/predicted)
  - Phenology
  - Range shifts (caveats)
- Coastal sage scrub in SD
  - Case studies: checkerspot butterfly, CA gnatcatcher
  - Response of CSS vs invasive plants to altered rainfall
  - Additional factors: Fire, N deposition
  - Restoration in the face of invasion and environmental change

#### **Coastal Sage Scrub**

- -One of the most endangered ecosystems in the US (Rubinoff 2001)
- $\approx$  200 species of forbs (Skinner & Pavlik 1994), many sensitive/listed under ESA
- $\approx$  100 listed/sensitive animal species in Southern CA CSS, 30+ birds, 30+ mammals, 25+ herps, several inverts

(www.rctlma.org/mshcp/volume1/index.html, www.sdcounty.ca.gov/pds/mscp/biology.html).

More than half has no protection (data from USGS 2013)



#### Quino checkerspot (Euphydryas editha quino)





Fig. 1. Quino checkerspot butterfly (QCB) study area and butterfly locations classified by colored circles according to the most recent decade in which an observation was reported. Also shown are locations of extinct (triangle) and extant (circle) populations used in modeling.


range, competition with

invasive plants

## California gnatcatcher response to climate change depends on CSS range shift




Climate change adaptation for many sensitive species will depend on reducing invasion by exotic annual grasses, which prevent establishment & growth by native CSS shrubs




Eliason & Allen 1997, Restoration Ecology

#### Outline



- Species responses to climate change (observed/predicted)
  - Phenology
  - Range shifts (caveats)
- Coastal sage scrub
  - Case studies: checkerspot butterfly, CA gnatcatcher
  - Response of CSS vs invasive plants to altered rainfall
  - Additional factors: Fire, N deposition
  - Restoration in the face of invasion and environmental change

## How will native CSS compete with exotic species with predicted climate change?



Diffenbaugh et al. 2008

- Coming decades are predicted to be warmer
   & drier
- Greater inter-annual variability in temperature & precipitation (extreme wet years and dry years)

#### Global Change Biology


Global Change Biology (2012) 18, 277-289, doi: 10.1111/j.1365-2486.2011.02480.x

#### Climate change and the invasion of California by grasses

BRODY SANDEL and EMILY M. DANGREMOND

Department of Integrative Biology, University of California, Berkeley, CA 94720, USA

- -Exotic species are currently found in warmer areas of California
- -Exotic annual grasses are particularly likely to have traits associated with warm areas
- -Predict exotic annual grasses will increase in abundance and extent with future warming
- -Patterns with rainfall are much less clear



## Do high rainfall years (or supplemental water) promote native shrub restoration?

- In past decades, waves of shrub establishment correspond with El Niño years
- Exotic annual species also often have high biomass under high resource supply
- Response of native shrubs to high rainfall may depend on competition with exotic species





#### Rainfall manipulation experiments

- UC Elliott Chaparral Reserve (MCAS Miramar), Santa Margarita Ecological Reserve (Fallbrook)
- Collect and reapply rainfall (drought to deluge)
- 3.5 x 3.5 m plots
- Measures:
  - Mature shrub growth
  - Growth and establishment of native shrub seedlings
  - Herbaceous species composition
     & production (mostly exotic)
  - Functional trait measures








Santa Margarita Ecological Reserve, Fallbrook

### Trade off: Native shrubs tolerate drought, exotic species dominate in high rainfall






- Restoration efforts involving watering, or in wet years need to pay particular attention to exotic species control
- Drought years offer windows of low competition for shrub establishment


Ashbacher et al. in review

# What source population to plant? Regional variation in *Artemisia californica* response to changing rainfall




- -Southern populations have experienced greater variation in yearly rainfall in the past
- -These populations respond most positively to increased watering






Pratt & Mooney 2013, Global Change Biology

#### Outline



- Species responses to climate change (observed/predicted)
  - Phenology
  - Range shifts (caveats)
- Coastal sage scrub
  - Case studies: checkerspot butterfly, CA gnatcatcher
  - Response of CSS vs invasive plants to altered rainfall
  - Additional factors: Fire, N deposition
  - Restoration in the face of invasion and environmental change

## Increasing fire frequency, positive feedbacks as vegetation responds to climate change



Westerling et al 2009, California Climate Change Center Draft Report "Climate change, growth, and California wildfire"

## Fire has direct effects on biota, promotes exotic invasion of CSS

- -Traditional fire-return intervals are hard to discern for CSS
- 20 years for Venturan CSS (Minnich & O'Leary 1986)
- 25 years in Riversidian CSS (Minnich unpub.)
- Fire regime has accelerated in many areas due to human activities, climate change, increase in exotic annual grasses due to N deposition



Recently burned hillside in Orange County, Photo credit: Jen Funk


# In SD county CSS, native shrub cover increases in the 20 years following fire




-Shrubs can recover following fire, even the presence of exotic species, given enough time

-Accelerating fire regimes may maintain exotic dominance

-Sites with high N deposition may see slow shrub declines






Gressard 2012, Dissertation, Data from USGS

#### **CSS & N Deposition**

- -Nitrogen depositionexceeds critical loads forCSS over much of SouthernCalifornia
- -Many studies show that exotic annual grasses benefit more than native species from N enrichment
- -Restoration practices should consider efforts to reduce N deposition (mulching), promote native species that can compete effectively with exotics



#### Outline



- Species responses to climate change (observed/predicted)
  - Phenology
  - Range shifts (caveats)
- Coastal sage scrub
  - Case studies: checkerspot butterfly, CA gnatcatcher
  - Response of CSS vs invasive plants to altered rainfall
  - Additional factors: Fire, N deposition
  - Restoration in the face of invasion and environmental change 2 case studies

- Case study: Laguna Coast
   Wilderness Park, Orange County CA
- Reserves at the urban-wildland interface have "edges" that are often along roads
- Invasive species, disturbance, N deposition
- Management along edges to maintain a buffer of native vegetation, to protect the interior
- Test strategies to reduce N availability and identify native species that can compete effectively with exotic species






## Hypothesis 1: Invasion will be reduced under lower N availability


#### Three levels of Nitrogen

**Ambient** 

- + N (9 g N/ $m^2$ yr)
- N (600 g  $C/m^2yr$ )







## Hypothesis 2: Native species prevent invasion by exotic species with similar functional traits

- Planted 6 seed mixes, each containing the same 20 native species, monitored invasion



-Annual grass (early)



-Early-active forb



-Perennial grass (late)



-Late-active forb



-Legume




-Even mix of all functional groups

#### Lowering N availability lowered invasion

- -Carbon addition strongly favored native over invasive species
- -Nitrogen enrichment leads to high dominance by exotic annual grasses








+ Carbon

+ Nitrogen


## Least cover of exotic species in plots dominated by early-active native forbs... why?



Cleland et al. 2012 Restoration Ecology

## Phenology – Exotic species have early phenology, compete most strongly with early-active native species

Aristida adcensionis Bromus madritensis Bromus diandrus Erodium cicutarium Lasthenia californica Lupinus succulentus Malva parviflora Anagalis arvensis Conyza floribunda Sonchus oleraceus Bromus hordeaceus Vulpia myuros Vulpia microstachys Lupinus bicolor Escholzia californica Lolium perene Phacelia cicutaria Raphanus sativus Medicago polymorpha Brassica nigra Hemizonia fasciculatum Nasella pulcrha Avena fatua Lolium multiflorum Vicia dasycarpa Clarkia purpurea Eriophylum confertiflorum Sisyrinchium bellum Bromus carinatus Nasella lepida Lotus purshianus



# Conclusion: Restoration through altered timing?

- Plant native species that have overlapping phenology with problematic invaders
- What allows exotic species to be active early in the growing season?
- Can we exploit this early phenology in restoration strategies?






# Exotic annual grasses have early phenology due to flexible germination cues


- Growth chamber study
  - 12 native spp, 12 exotic spp
  - Cues: soil moisture, daylength, temperature
  - Exotic species were better able to maximize germination under favorable growing conditions





#### Benefits and costs of early phenology


- Exotic species could benefit from seasonal priority effects
  - germinate earlier,
     preempt resources,
     competitively dominant




- Costs of early phenology
  - Risk of missing the true start of the growing season
  - Apparency to herbivores




Hyp 1: Priority advantage: flexible germination cues



Hyp 1: Priority advantage: flexible germination cues




Hyp 2: Priority disadvantage: mistiming





Case study 2: Timed watering to favor native over exotic species

- UC Scripps Coastal Reserve
- Compared plots with late-summer watering to unwatered controls (winter rains only)
- Focal native & exotic species were seeded into each plot, in addition to existing seedbank
- Monitored germination and survivorship












#### A. Winter Rains, priority advantage




#### A. Winter Rains, priority advantage







#### **Conclusions**

- SDMs provide an important starting point for predicting species at risk for declines with climate change
- Lots of variability... success is likely based on monitoring populations & evaluating strategies prior to implementation
- Strategies for restoration & adaptation to climate change need to consider other interacting environmental factors
  - Shifting habitat, species interactions
  - Invasive species
  - N deposition
  - Fire regimes
- Opportunities for creative & effective conservation of sensitive species and habitat









#### Talk outline



#### 1. What is Climate Adaptation?

- 1. General principles
- 2. Aligning with management goals

#### 2. Adaptation Planning

- 1. Frameworks
- 2. An example from Yellowstone
- 3. In the context of INRMPs & Det. Fallbrook



3. Objectives of breakout groups

## Climate adaptation



Actions that enable species, ecosystems—and humans—better cope with or adjust to changing conditions.









## Climate adaptation



Conceptually challenging

High Uncertainty





High Complexity

## General principles



- Reduce non-climate stressors
- Manage for ecological function and biodiversity
- Establish buffer zones and connectivity
- Increase monitoring & facilitate management under uncertainty
- Implement "proactive" strategies



Adapted from "Adaptation 2009" discussion paper by Glick et al. and other sources

## Aligning management goals




### What are they trying to achieve?

- Increase **RESISTANCE** to change?
- Promote RESILIENCE to change?
- To enable ecosystems to RESPOND to change? (sometimes called facilitation)
- To help determine when to consider TRIAGE?




## Goal of adaptation planning





### Adaptation framework






**Step 2: Assess effects on target** 

## An adaptation framework







Scenario 1

Scenario 2

Scenario 3

#### **3b. MANAGEMENT RESPONSES**

Management Action 1

Management Action 2

Management Action 3

Management Action 4

#### 3a. EFFECTS ON TARGET

Scenario 1

Scenario 2

Scenario 3

#### THE ACT FRAMEWORK Implementation & **Adaptation planning phase** efficacy evaluation 1. Select conservation target phase and Define management objective 2. Assess 6. Monitor and climate change effects evaluate action plan efficacy **Build** Repeat for: conceptual more targets model more 6.5 Adjust actions Identify objectives Assess as needed future Responses more climate to scenarios scenarios information 5. Implement priority actions 3. Identify intervention points and management actions Cross et al. 2012, 4. Prioritize actions &

**Env Mangmnt** 

develop plan

### An example



Yellowstone River flows (ecological process)

# Step 1: select target & objective




To maintain Yellowstone cutthroat trout



### An example





Step2: Build conceptual model


### An example






## An example






## An example





#### Step 3b. Potential actions for initial climate scenario




#### Context of INRMPs



- Compliance with applicable natural resource laws (including regulations)
- Sustainable management of natural resources.
- Sustainable use by the public of resources subject to mission compatibility, safety & security
- NO NET LOSS in the capability of installation lands to support the military mission



# Detachment Fallbrook







# Targets for this workshop




#### 1. Coastal sage scrub



## **Breakout groups**






# Day One objectives

Day Two objectives

## Next steps



# Towards strategic "climate-informed" monitoring as a key adaptation action









# "Adapt, improvise, and overcome."

- T.C. Cummings, Mind of a SEAL




# IMPLANTING AT DETACHMENT



OR.. HOW IS THIS GOING TO WORK?

#### **HOW ARE WE TO PROCEED???**



#### LUNCH!!!



#### LUNCH.



#### **BREAK INTO GROUPS**



| Time   | Topic                       | Group Facilitators             |
|--------|-----------------------------|--------------------------------|
| 12:30- | Group 1: Coastal Sage Scrub | Robert Wolf & Brenna Vredeveld |
| 16:00  |                             |                                |
| 12:30- | Group 2: Riparian Woodland  | Bill Wild & Anna Kellogg       |
| 16:00  |                             |                                |

#### DAY 1, STEP 1 & 2

1. What is our targeted conservation feature?



**Coastal Sage Scrub** 



**Riparian Woodlands** 

2. Review the current INRMP management objectives for CSS or Riparian.

#### DAY 1, STEP 3

Conceptual model review, revision.





#### DAY 1, STEP 4

Use the conceptual model as a guide to filling in the blanks in Table 1.

Table 1: Primary reference for linking Climate Change with our hypotheses of ecological change

Identify Key Driver - How does the driver manifests under different scenarios? - What is our hypothesis of ecological change on our conservation target?

Finally – what monitoring activities are relevant to the ecological hypotheses we come up with?

### DAY 1, STEP 5

Let it sink in.



Sleep on it.

#### DAY 2, ALL MORNING

Day two is all Management Focused.

Start with our Hypotheses of Change from Table 1. and use it to begin Table 2.

This will allow us think systematically about how we monitor (Drivers and Pressures as well as our hypothesized effects).

Management under different scenarios.

The priorities of our proposed interventions.

Opportunities for implementation.

Re-evaluation of management objectives.

#### DAY 2, AFTERNOON - REVIEW AND SYNTHESIS

#### Part 1. You present your work!

Identify your strategic actions, and their rank.

General thoughts, considerations, concerns regarding management actions.

#### Part 2. Facilitated discussion by Carolyn Enquist.

What barriers might exist for implementing strategic actions?

How to overcome them?

Installation versus regional action?

#### **NEXT STEPS AND WRAP-UP**


If monitoring is so important, how are going to set the stage to implement it? – Carolyn Enquist


Workshop Summary and what are going to do with all this information? – Dawn Lawson

Goodbye. ⊗ - Christy Wolf.

#### **LETS EAT!**








Carolyn Enquist, Science Coordinator USA-NPN Det. Fallbrook Adaptation Planning Workshop 28-29 August 2013



# Climate adaptation









# "Adapt, improvise, and overcome."

- T.C. Cummings, Mind of a SEAL

# Goal of adaptation planning





# Targets & objectives




1. Coastal sage scrub: maintain habitat for gnatcatchers & SKR



## **Breakout groups**





Day One objectives

Day Two objectives

#### Step 3b. Potential actions for initial climate scenario

Potential Actions -Intervention Points Desired Responses Riparian **Conduct & Implement** Restoration, Identify Maximize water management Fluvial & veg risk assessment infrastructure availability Improvement, repair, Fire reduce channelization Exclusion fencing, shading management **Improve** Reduce **Exotic spp** Implement eradication tools Suitable habitat: competition & habitat management degradation; Targeted herbicide treatments Retain and maximize buffer neg effects water in system fm fire Balance needs of Protection of **Prevention & suppression** SKR & gnatcatcher Refugia & measures Grazing seed sources Reduce biomass of annual exotics

#### Discussion



# How would you describe priority actions?

- RESISTANCE
- RESILIENCE
- RESPONSE or facilitation
- RE-ALIGNMENT
- TRIAGE?



Adapted from Millar et al. 2007, Ecological Applications and USFS Climate Change Resource Center

#### Context of INRMPs



- Compliance with applicable natural resource laws (including regulations)
- Sustainable management of natural resources.
- Sustainable use by the public of resources subject to mission compatibility, safety & security
- NO NET LOSS in the capability of installation lands to support the military mission



#### Context of INRMPs




#### Detachment Fallbrook


- Barriers to implementation?
- Opportunities?


## Next topic



# Towards strategic "climate-informed" monitoring as a key adaptation action







Carolyn Enquist, Science Coordinator USA-NPN Det. Fallbrook Adaptation Planning Workshop 28-29 August 2013





#### C-I monitoring?



C-I (or climate-smart) monitoring helps provide context for understanding climate-related impacts and vulnerabilities and for informing agile adaptive management.

Monitoring approaches should be carefully designed to ensure they are capable of guiding needed adjustments in strategies and actions—evaluation.

--Stein et al. "ClimateSmart Conservation Quick Guide" (full document in prep.)

#### What to monitor?



### Essential Biodiversity Variables (EBVs)—Pereira et al. 2013, *Science*

| EBV CLASS               | EBV EXAMPLES                          |
|-------------------------|---------------------------------------|
| (1) Genetic composition | Allelic diversity, phenotypes         |
| (2) Species populations | Abundance & distribution              |
| (3) Species traits      | Phenology                             |
| (4) Composition         | Taxonomic diversity, functional types |
| (5) Ecosystem structure | Habitat structure                     |
| (6) Ecosystem function  | Ecosystem processes                   |

#### Other considerations



#### **Motivation**

- Status & trends (discovery)
- Question driven (evidence)
- Indicators of change
- Funding

#### **Design:**

- Timing, Frequency
- Co-location of weather stations, soil moisture, another biodiversity metric?

#### **Data Management & Analysis:**

- Storage
- Analysis










## The role of Phenology in strategic Climate-Informed Monitoring

Carolyn Enquist, Science Coordinator USA-NPN Det. Fallbrook Adaptation Planning Workshop 28-29 August 2013





#### Observed shifts in the SW region



- Shift in timing of flowering (Bowers 2007, Inouye 2008, Kimball et al. 2009, Crimmins et al. 2011)
- Timing of migratory bird arrivals

(McKinney et al. 2012, Kellermann in prep)

- Timing of animal emergence (Inouye et al. 2000, Forister & Shapiro 2003, Ozgul et al. 2010)
- Timing of egg laying (Brown et al. 1999)



#### What to monitor?

## Essential Biodiversity Variables (EBVs)—Pereira et al. 2013, *Science*

| EBV CLASS               | EBV EXAMPLES                           |
|-------------------------|--|
| (1) Genetic composition | Allelic diversity, phenotypes          |
| (2) Species populations | Abundance & distribution               |
| (3) Species traits      | Phenology                              |
| (4) Composition         | Taxonornic diversity, functional types |
| (5) Ecosystem structure | Habitat structure                      |
| (6) Ecosystem function  | Ecosystem processes                    |

Spatial Extent

Seasonal

Annual

Decadal

#### **Temporal Scale**



#### **Ecological Level**

Species -Individuals

Community-Population Biomes -Ecosystems

#### **Continents to Globe:**

- Growing degree days
- Timing and duration of freeze events
- Timing & frequency of extreme events
- Synchronicity and natural variation

#### **Landscapes to Regions:**

- Timing of snowmelt & peak streamflows
- Start of season, duration & end of season
- Start of fire season & duration

#### **Species to Landscapes:**

Timing of


- leaf and bloom
- animal emergence & migrations




# Key indicator of climate change impacts on ecosystems:

"Phenology...is perhaps the simplest process in which to track changes in the ecology of species in response to climate change."

(IPCC 2007)






#### **Applications**



### Predicting species response & vulnerability: Implications for

invasiveness, sensitivity & adaptive capacity



Change in phenology





Willis et al. 2008 PNAS Moller et al. 2008 PNAS Willis et al. 2010 PLOS Biology Ozgul et al. 2010 Nature Hulme 2011 New Phyt. Cleland et al. 2012


#### **Applications**



#### **Shifts in the Timing of Management Actions**



#### Supporting decision-making



## Toward a framework for strategic climate-informed monitoring for INRMPs....

Can we come up with a shared vision?

#### Other considerations



#### **Motivation**

- Status & trends
- Question driven
- Funding

#### **Design:**

- Timing
- Frequency
- Co-location of weather stations, soil moisture, another biodiversity metric?

#### **Data Management & Analysis:**

- Storage
- Analysis



